

INFINITE DIMENSIONAL GEOMETRY OF $M_1 = \text{Diff}_+(\mathbb{S}^1)/\text{PSL}(2, \mathbb{R})$ AND q_R -CONFORMAL SYMMETRIES

DENIS V. JURIEV

June, 24, 1988

math.RT/9806140

This article being addressed to specialists in representation theory, quantum algebra, infinite dimensional differential (Kähler) geometry, mathematical physics and informatics is devoted to various objects of infinite dimensional geometry related to q_R -conformal symmetries, which analysis being initiated by the author primarily for goals of informatics of the interactive videosystems acquired motivations from pure mathematics later. On the other hand the constructions considered below are originated ideologically in a certain extent in such branches of modern mathematical physics as conformal and string theories and these origins penetrated deeply into the view on nature of the investigated objects. Precisely, the interrelations of the infinite dimensional geometry of the homogeneous Kähler manifold $M = \text{Diff}_+(\mathbb{S}^1)/\mathbb{S}^1$ [1,2] (which plays a crucial role in the geometric formulation of conformal and string field theories [3-5]), its quotient $M_1 = \text{Diff}_+(\mathbb{S}^1)/\text{PSL}(2, \mathbb{R})$ and the approximate representations of the Virasoro algebra by the q_R -conformal symmetries [6] are discussed. I believe that the obtained results may strengthen and clarify the connections between infinite dimensional geometry, representation theory, modern mathematical physics and informatics of interactive systems once suggested in [2,4:III].

1. PRELIMINARY DEFINITIONS

1.1. The Lie algebra $\mathfrak{sl}(2, \mathbb{C})$ and Verma modules over it. Lie algebra $\mathfrak{sl}(2, \mathbb{C})$ is a three-dimensional space of 2×2 complex matrices with zero trace supplied by the standard commutator $[X, Y] = XY - YX$, where the right hand side multiplication is the standard matrix multiplication. In the basis

$$l_{-1} = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}, \quad l_0 = \begin{pmatrix} \frac{1}{2} & 0 \\ 0 & -\frac{1}{2} \end{pmatrix}, \quad l_1 = \begin{pmatrix} 0 & 0 \\ -1 & 0 \end{pmatrix}$$

the commutation relations are of the form:

$$[l_i, l_j] = (i - j)l_{i+j} \quad (i, j = -1, 0, 1).$$

Lie algebra $\mathfrak{sl}(2, \mathbb{C})$ is $\mathbb{Z}$ -graded: $\deg(l_i) = -\text{ad}(l_0)l_i = i$, where $\text{ad}(X)$ is the adjoint action operator in the Lie algebra: $\text{ad}(X)Y = [X, Y]$. Therefore, $\mathbb{Z}$ -graded modules over $\mathfrak{sl}(2, \mathbb{C})$ are l_0 -diagonal. A vector v in a $\mathbb{Z}$ -graded module over the Lie algebra $\mathfrak{sl}(2, \mathbb{C})$ is called extremal iff $l_1 v = 0$ and the linear span of vectors $l_{-1}^n v$

($n \in \mathbb{Z}_+$) coincides with the module itself (i.e. v is a cyclic vector). A $\mathbb{Z}$ -graded module with an extremal vector (in this case it is defined up to a multiplier) is called extremal [7]. An extremal module is called the Verma module [8] iff the action of l_{-1} is free in it, i.e. the vectors $l_{-1}^n v$ are linearly independent. In the case of the Lie algebra $\mathfrak{sl}(2, \mathbb{C})$ the Verma modules are just the same as infinite dimensional extremal modules. An extremal weight of the Verma module is the number defined by the equality $l_0 v = h v$, where v is the extremal vector. The Verma modules are defined for all complex numbers h and are pairwise nonisomorphic. Below we shall consider the Verma modules with real extremal weights only.

The Verma module V_h over the Lie algebra $\mathfrak{sl}(2, \mathbb{C})$ with the extremal weight h may be realized in the space $\mathbb{C}[z]$ of polynomials of a complex variable z . The formulas for the generators of the Lie algebra $\mathfrak{sl}(2, \mathbb{C})$ are of the form:

$$l_{-1} = z, \quad l_0 = z \partial_z + h, \quad l_1 = z \partial_z^2 + 2h \partial_z,$$

here $\partial_z = \frac{d}{dz}$.

The Verma module is nondegenerate (i.e. does not contain any proper submodule) iff $h \neq -\frac{n}{2}$ ($n \in \mathbb{Z}_+$). The Verma module V_h is called unitarizable (or hermitean) iff it admits a structure of the pre-Hilbert space such that $l_i^* = l_{-i}$. The completion of the unitarizable Verma module will be denoted by V_h^{Hilb} . The Lie algebra $\mathfrak{sl}(2, \mathbb{C})$ acts in V_h^{Hilb} by the unbounded operators. Also it is rather useful to consider the formal Verma modules V_h^{form} , which are realized in the space $\mathbb{C}[[z]]$ of formal power series of a complex variable z , whereas the formulas for generators of the Lie algebra $\mathfrak{sl}(2, \mathbb{C})$ coincide with ones above. Note that $V_h \subseteq V_h^{\text{Hilb}} \subseteq V_h^{\text{form}}$ and modules $V_h, V_h^{\text{Hilb}}, V_h^{\text{form}}$ form the Gelfand triple or the Dirac equipment of the Hilbert space V_h^{Hilb} . An action of the real form of the Lie algebra $\mathfrak{sl}(2, \mathbb{C})$ generated by the anti-Hermitean operators $i l_0, l_1 - l_{-1}$ and $i(l_1 + l_{-1})$ in the Hilbert space V_h^{Hilb} by the unbounded operators is exponentiated to a unitary representation of the corresponding simply connected Lie group.

In the nonunitarizable Verma module over the Lie algebra $\mathfrak{sl}(2, \mathbb{C})$ there exists the unique (up to a scalar multiple) indefinite sesquilinear form $(\cdot, \cdot)$ such that $(l_i v_1, v_2) = (v_1, l_{-i} v_2)$ for any two vectors v_1 and v_2 from the Verma module. If this sesquilinear form is nondegenerate (in this case the Verma module is nondegenerate) then it has a signature (n, ∞) , where n is finite, and therefore, there is defined a Pontryagin completion [9] of the Verma module. The corresponding module in which the Lie algebra $\mathfrak{sl}(2, \mathbb{C})$ acts by the unbounded operators will be denoted by V_h^{Pontr} . The following chain of inclusions holds: $V_h \subseteq V_h^{\text{Pontr}} \subseteq V_h^{\text{form}}$. An action of the real form of the Lie algebra $\mathfrak{sl}(2, \mathbb{C})$ generated by the anti-Hermitean (with respect to the nondegenerate indefinite sesquilinear form $(\cdot, \cdot)$) operators $i l_0, l_1 - l_{-1}$ and $i(l_1 + l_{-1})$ in the Pontryagin space V_h^{Pontr} by the unbounded operators is exponentiated to a pseudounitary representation of the corresponding simply connected Lie group.

1.2. Hidden symmetries in Verma modules over the Lie algebra $\mathfrak{sl}(2, \mathbb{C})$: Lobachevskii-Berezin C^* -algebra and q_R -conformal symmetries.

Proposition 1 [10]. *In the nondegenerate Verma module V_h over the Lie algebra $\mathfrak{sl}(2, \mathbb{C})$ there are uniquely defined the operators D and F such that*

$$\begin{aligned} [D, l_{-1}] &= 1, & [D, l_0] &= D, & [D, l_1] &= D^2, \\ [l_{-1}, F] &= 1, & [l_0, F] &= F, & [l_1, F] &= F^2. \end{aligned}$$

If the Verma modules are realized in the space $\mathbb{C}[z]$ of polynomials of a complex variable z then

$$D = \partial_z, \quad F = z \frac{1}{\xi + 2h},$$

where $\xi = z\partial_z$. The operators F and D obey the following relations:

$$[FD, DF] = 0, \quad [D, F] = q_R(1 - DF)(1 - FD),$$

where $q_R = \frac{1}{2h-1}$. In the unitarizable Verma module ($q_R \neq 0$) the operators F and D are bounded and $F^* = D$, $D^* = F$.

The algebra generated by the variables t and t^* with the relations $[tt^*, t^*t] = 0$ and $[t, t^*] = q_R(1 - tt^*)(1 - t^*t)$ being the Berezin quantization of the Lobachevskii plane realized in the unit complex disc (the Poincaré realization) [11] is called the Lobachevskii-Berezin algebra. Proposition 1 allows to consider the Lobachevskii-Berezin algebra as a C^* -algebra. The Lobachevskii-Berezin C^* -algebra was rediscovered recently by S.Klimek and A.Lesniewsky [12].

Proposition 2 [10]. *In the nongenerate Verma module V_h over the Lie algebra $\mathfrak{sl}(2, \mathbb{C})$ there are uniquely defined the operators L_n ($n \in \mathbb{Z}$) such that*

$$[l_i, L_n] = (i - n)L_{i+n} \quad (i = -1, 0, 1; n \in \mathbb{Z}),$$

moreover, $L_i = l_i$ ($i = -1, 0, 1$). If the Verma modules are realized in the space $\mathbb{C}[z]$ of polynomials of a complex variable z then

$$L_k = (xi + (k+1)h)\partial_z^k \quad (k \geq 0), \quad L_{-k} = z^k \frac{\xi + (k+1)h}{(\xi + 2h) \dots (\xi + 2h + k - 1)} \quad (k \geq 1),$$

where $\xi = z\partial_z$. The operators L_n obey the following relations:

$$[L_n, L_m] = (n - m)L_{n+m}, \quad \text{if } n, m \geq -1 \text{ or } n, m \leq 1.$$

In the unitarizable Verma module the operators L_n are unbounded and $L_i^* = L_{-1}$.

The operators L_n are called the q_R -conformal symmetries. They may be symbolically represented in the form:

$$L_n = D^{nh} L_0 D^{n(1-h)}, \quad L_{-n} = F^{n(1-h)} L_0 F^{nh}.$$

To supply the symbolical recording by a sense one should use the general commutation relations

$$[L_n, f(D)] = (-D)^{n+1} f'(D) \quad (n \geq -1), \quad [L_{-n}, f(F)] = F^{n+1} f'(D) \quad (n \geq -1)$$

for $n = 0$.

The commutation relations for the operators D , F and the generators of q_R -conformal symmetries with the generators of the Lie algebra $\mathfrak{sl}(2, \mathbb{C})$ mean that the families J_k and L_k ($k \in \mathbb{Z}$), where $J_i = D^i$, $J_{-i} = F^i$ ($i \in \mathbb{Z}_+$), are families of tensor operators [9,13] for the Lie algebra $\mathfrak{sl}(2, \mathbb{C})$.

1.3. Infinite dimensional $\mathbb{Z}$ -graded Lie algebras: the Witt algebra $\mathfrak{w}^{\mathbb{C}}$ of all Laurent polynomial vector fields on a circle and the Virasoro algebra $\text{vir}^{\mathbb{C}}$, its one-dimensional nontrivial central extension. The Lie algebra $\text{Vect}(S^1)$ is realized in the space of C^∞ -smooth vector fields $v(t)\partial_t$ on a circle $S^1 \simeq \mathbb{R}/2\pi\mathbb{Z}$ with the commutator

$$[v_1(t)\partial_t, v_2(t)\partial_t] = (v_1(t)v_2'(t) - v_1'(t)v_2(t))\partial_t.$$

In the basis

$$s_n = \sin(nt)\partial_t, \quad c_n = \cos(nt)\partial_t, \quad h = \partial_t$$

the commutation relations have the form:

$$\begin{aligned} [s_n, s_m] &= \frac{1}{2}((m-n)s_{n+m} + \text{sgn}(n-m)(n+m)s_{|n-m|}), \\ [c_n, c_m] &= \frac{1}{2}((n-m)s_{n+m} + \text{sgn}(n-m)(n+m)s_{|n-m|}), \\ [s_n, c_m] &= \frac{1}{2}((m-n)c_{n+m} - (m+n)c_{|n-m|}) - n\delta_{nm}h, \\ [h, s_n] &= nc_n, \quad [h, c_n] = -ns_n. \end{aligned}$$

Let us denote by $\text{Vect}^{\mathbb{C}}(S^1)$ the complexification of the Lie algebra $\text{Vect}(S^1)$. In the basis $e_n = ie^{ikt}\partial_t$ the commutation relations in the Lie algebra $\text{Li Vect}^{\mathbb{C}}(S^1)$ have the form:

$$[e_j, e_k] = (j-k)e_{j+k}.$$

It is rather convenient to consider an imbedding of the circle S^1 into the complex plane $\mathbb{C}$ with the coordinate z , so that $z = e^{it}$ on the circle and the elements of the basis e_k ($k \in \mathbb{Z}$) are represented by the Laurent polynomial vector fields:

$$e_k = z^{k+1}\partial_z.$$

The $\mathbb{Z}$ -graded Lie algebra generated by the Laurent polynomial vector fields (i.e. by the finite linear combinations of elements of the basis e_k) is called the Witt algebra and is denoted by $\mathfrak{w}^{\mathbb{C}}$. The Witt algebra $\mathfrak{w}^{\mathbb{C}}$ is the complexification of the subalgebra $\mathfrak{w}$ of the algebra $\text{Vect}(S^1)$ generated by the trigonometric polynomial vector fields on a circle S^1 , i.e. by the finite linear combinations of elements of the basis s_n, c_n and h .

The Lie algebra $\text{Vect}(S^1)$ admits a nontrivial one-dimensional central extension defined by the Gelfand-Fuchs 2-cocycle [14]:

$$c(v_1(t)\partial_t, v_2(t)\partial_t) = \int_0^{2\pi} (v_1'(t)v_2''(t) - v_2'(t)v_1''(t)) dt.$$

This extension being continued to the complexification $\text{Vect}^{\mathbb{C}}(S^1)$ of the Lie algebra $\text{Vect}(S^1)$ and reduced to the subalgebra $\mathfrak{w}^{\mathbb{C}}$ defines a central one-dimensional extension of the Witt algebra, which is called the Virasoro algebra and is denoted by $\text{vir}^{\mathbb{C}}$. The Virasoro algebra is generated by the elements e_k ($k \in \mathbb{Z}$) and the central element c with the commutation relations:

$$[e_j, e_k] = (j-k)e_{j+k} + \frac{j^3-j}{12}c\delta(j+k)$$

and is the complexification of a central extension vir of the Lie algebra $\mathfrak{w}$. Because the 2-cocycle

$$\int_0^{2\pi} (v_1(t)v_2'(t) - v_2(t)v_1'(t)) dt$$

is trivial, the cocycle, which defines the Virasoro algebra, is indeed equivalent to the Gelfand-Fuchs 2-cocycle above and is known under the same name. The modified Gelfand-Fuchs cocycle is $\mathfrak{sl}(2, \mathbb{C})$ -invariant, where $\mathfrak{sl}(2, \mathbb{C})$ is a subalgebra of $\text{vir}^{\mathbb{C}}$ generated by e_{-1} , e_0 and e_1 , so it is handier in practice. Below we shall use the modified version of Gelfand-Fuchs 2-cocycle under this name only. In the irreducible representation the central element c of the Virasoro algebra is mapped to a scalar operator, which is proportional to the identity operator with a coefficient called *the central charge*.

2. $\mathcal{HS}$ - AND $\mathcal{TC}$ -PROJECTIVE REPRESENTATIONS OF THE WITT ALGEBRA

2.1. $\mathfrak{A}$ -projective representations [6].

Definition 1 [6].

A Let $\mathfrak{A}$ be an associative algebra represented by linear operators in the linear space H and $\mathfrak{g}$ be the Lie algebra. The linear mapping $T : \mathfrak{g} \mapsto \text{End}(H)$ is called a *$\mathfrak{A}$ -projective representation* iff for any X and Y from $\mathfrak{g}$ there exists an element of the algebra $\mathfrak{A}$ represented by the operator A_{XY} such that

$$[T(X), T(Y)] - T([X, Y]) = A_{XY}.$$

The operators A_{XY} are called *deviations* for the $\mathfrak{A}$ -projective representation T .

B Let $\mathfrak{A}$ be an associative algebra with an involution $*$ symmetrically represented in the Hilbert space H . If $\mathfrak{g}$ is a Lie algebra with an involution $*$ then its $\mathfrak{A}$ -projective representation T in the space H is called *symmetric* iff for all elements a from $\mathfrak{g}$ $T(a^*) = T^*(a)$. Let $\mathfrak{g}$ be $\mathbb{Z}$ -graded Lie algebra ($\mathfrak{g} = \bigoplus_{n \in \mathbb{Z}} \mathfrak{g}_n$) with an involution $*$ such that $\mathfrak{g}_n^* = \mathfrak{g}_{-n}$ and involution is identical on the subalgebra $\mathfrak{g}_0$. Let us extend the $\mathbb{Z}$ -grading and the involution $*$ from the Lie algebra $\mathfrak{g}$ to the tensor algebra $\mathbf{T}(\mathfrak{g})$. A symmetric $\mathfrak{A}$ -projective representation of $\mathfrak{g}$ is called *absolutely symmetric* iff for any element a of the algebra $\mathbf{T}(\mathfrak{g})$ such that $\deg(a) = 0$ the identity $T(a) = T^*(a)$ holds (here the representation T of the Lie algebra $\mathfrak{g}$ in H is extended to the mapping of $\mathbf{T}(\mathfrak{g})$ into $\text{End}(H)$).

C A $\mathfrak{A}$ -projective representation T of the Lie algebra $\mathfrak{g}$ in the linear space H is called *almost absolutely closed* iff for any natural n and arbitrary elements $X_0, X_1, X_2, \dots, X_{n+1}$ of the Lie algebra $\mathfrak{g}$ there exists an element $\varphi(X_0, X_1, X_2, \dots, X_{n+1})$ of the algebra $\mathfrak{g}$ such that

$$[\dots [[T(X_0), T(X_1)], T(X_2)], \dots, T(X_{n+1})] \equiv T(\varphi(X_0, X_1, X_2, \dots, X_{n+1})) \pmod{\mathfrak{A}},$$

here $\mathfrak{A}$ is considered as represented in $\text{End}(H)$. An almost absolutely closed $\mathfrak{A}$ -projective representation T of the Lie algebra $\mathfrak{g}$ in the linear space H is called *absolutely closed* iff $\varphi(\cdot, \dots, \cdot) \equiv 0$.

If H is infinite dimensional then the representation may be realized by the unbounded operators with suitable domains of definition.

Remark 1. The standard projective representation is a particular case of Def.1A if the algebra $\mathfrak{A}$ is one-dimensional and acts in H by scalar operators.

Remark 2. If H is a Hilbert (or pre-Hilbert) space then one may consider the algebra $\mathcal{HS}$ of all Hilbert-Schmidt operators as $\mathfrak{A}$. It is possible to consider the algebras $\mathcal{B}$, $\mathcal{K}$, $\mathcal{TC}$ and $\mathcal{FR}$ of all bounded, compact, trace class and finite-rank operators, too.

Remark 3. One may define $\mathfrak{a}$ -projective representations of Lie algebra $\mathfrak{g}$, where $\mathfrak{a}$ is a Lie algebra in the similar way. Any $\mathfrak{A}$ -projective representation, where $\mathfrak{A}$ is an associative algebra, is a $\mathfrak{A}_{[\cdot,\cdot]}$ -projective representation, where $\mathfrak{A}_{[\cdot,\cdot]}$ is a commutator Lie algebra of the associative algebra $\mathfrak{A}$.

2.2. $\mathcal{HS}$ - and $\mathcal{TC}$ -projective representations of Lie algebras. Below we shall consider either $\mathcal{HS}$ - or $\mathcal{TC}$ -projective absolutely closed representations of Lie algebras only. Let's introduce some useful concepts.

Definition 2A. Let T be a $\mathcal{TC}$ -projective representation of Lie algebra $\mathfrak{g}$. Then the *fundamental form* α_{XY} on $\mathfrak{g}$ for the representation T will be defined as

$$\alpha_{XY} = \text{Tr } A_{XY},$$

where A_{XY} are deviations for representation T , $X, Y \in \mathfrak{g}$.

Let us denote by $\text{Ker } A = \{X \in \mathfrak{g} : (\forall Y \in \mathfrak{g}) A_{XY} = 0\}$ and $\text{Ker } \alpha = \{X \in \mathfrak{g} : (\forall Y \in \mathfrak{g}) \alpha_{XY} = 0\}$; $\mathfrak{g}_0 = \text{Ker } A$ is a Lie subalgebra of $\mathfrak{g}$. Note that the fundamental form α is $\mathfrak{g}_0$ -invariant. Let us suppose that $\text{Ker } A = \text{Ker } \alpha$ then one can define the formal series

$$A = \alpha^{XY} A_{XY}$$

(here and below, the Einstein rule of summation is assumed). If it converges absolutely in certain topology in some class $\mathcal{O}$ of operators, the result will be called *the mean deviation*.

Remark 4. The operator A is $\mathfrak{g}_0$ -invariant.

If T is (operator) irreducible representation of $\mathfrak{g}_0$ then one can define the *deviation constant* α as

$$A = \alpha E,$$

where E is the identical operator (note that α is not obligatory equal to 1 in the considered infinite-dimensional case).

Let us also define $B_{XYZ} = [T(X), A_{YZ}]$ and $\beta_{XYZ} = \text{Tr } B_{XYZ}$ (the least expressions exist because T is absolutely closed).

Remark 5.

$$d\alpha_{XYZ} + (\beta_{XYZ} + \beta_{YZX} + \beta_{ZXY}) = 0,$$

where

$$d\alpha_{XYZ} = \alpha_{X,[Y,Z]} + \alpha_{Y,[Z,X]} + \alpha_{Z,[X,Y]}.$$

The equalities follow from the identity

$$(A_{X,[Y,Z]} + B_{XYZ}) + (A_{Y,[Z,X]} + B_{YZX}) + (A_{Z,[X,Y]} + B_{ZXY}) = 0.$$

Definition 2B. Let T be a $\mathcal{HS}$ -projective representation of Lie algebra $\mathfrak{g}$. Then the *curvature form* $R \in S^2\Lambda^2(\mathfrak{g}^*)$ of T is defined as

$$R_{XY,UV} = \text{Tr}(A_{XY}A_{UV}).$$

If T is also $\mathcal{TC}$ -projective representation and $\text{Ker } A = \ker \alpha$ then let us introduce the *curvature operator*

$$R_{XY,U}^V = \alpha^{ZV} R_{XY,UZ}.$$

If the series $R_{XV,Y}^V$ and $R_{XY,V}^V$ absolutely converges they will be called the *Ricci tensors* of $\mathcal{TC}$ -projective representation T of the first and the second kind, respectively, and denoted by $R_{XY}^{(i)}$ ($i = 1, 2$). The $\mathcal{TC}$ -projective representation T will be called *einsteinian* (of the 1st or 2nd kind) iff $R_{XY}^{(i)}$ is proportional to the fundamental form α :

$$R_{XY}^{(i)} = \varrho^{(i)} \alpha_{XY}.$$

Such definition of Ricci tensors is motivated by analogies from Kähler geometry, where Ricci tensor may be defined equivalently via curvature operator as $R_{XV,Y}^V$ or as $R_{XY,V}^V$ [15].

Remark 6. Let T be a $\mathcal{TC}$ -projective representation of Lie algebra $\mathfrak{g}$, which is an (operator) irreducible representation of $\mathfrak{g}_0$ (and $\text{Ker } A = \text{Ker } \alpha$), and the formal series for the mean deviation A absolutely converges then the series for $R_{XY}^{(2)}$ also converges and $\varrho^{(2)} = \alpha$.

2.3. $\mathcal{HS}$ -projective representation of the Witt algebra by q_R -conformal symmetries in the unitarizable Verma modules V_h over the Lie algebra $\mathfrak{sl}(2, \mathbb{C})$. Note that the Witt algebra $\mathfrak{w}^{\mathbb{C}}$ admits a natural involution $*$.

Proposition 3 [6]. *The generators L_k ($k \in \mathbb{Z}$) of q_R -conformal symmetries in the unitarizable Verma module V_h over the Lie algebra $\mathfrak{sl}(2, \mathbb{C})$ realize an absolutely symmetric $\mathcal{HS}$ -projective representation of the Witt algebra $\mathfrak{w}^{\mathbb{C}}$.*

Remark 7. The $\mathcal{HS}$ -projective representations of the Witt algebra in the unitarizable Verma modules over $\mathfrak{sl}(2, \mathbb{C})$ are absolutely closed.

Remark 8. Proposition 3 is generalized for the pseudo-unitary case with the substitution of the class $\mathcal{HS}$ of Hilbert-Schmidt operators to the class $\mathcal{K}$ of compact operators.

Theorem 1. *The $\mathcal{HS}$ -projective representations of the Witt algebra $\mathfrak{w}^{\mathbb{C}}$ in the unitarizable Verma modules V_h over the Lie algebra $\mathfrak{sl}(2, \mathbb{C})$ by q_R -conformal symmetries are $\mathcal{TC}$ -projective representations. The fundamental form α_{XY} in the basis e_k has the form*

$$\alpha_{ij} = \frac{6h^2 - 6h + 1}{6}(j^3 - j)\delta(i + j).$$

Proof. In the considered case $\text{Ker } A = \mathfrak{sl}(2, \mathbb{C})$, hence the fundamental form α_{XY} is $\mathfrak{sl}(2, \mathbb{C})$ -invariant. All $\mathfrak{sl}(2, \mathbb{C})$ -invariant 2-forms on the Witt algebra $\mathfrak{w}^{\mathbb{C}}$ are proportional to the Gelfand-Fuchs cocycle. To determine the multiplier it is sufficient to calculate $\alpha_{2,-2}$ using explicit formulas for q_R -conformal symmetries.

Remark 9. If $h = \frac{1}{2} \pm \frac{\sqrt{3}}{6}$ then $\alpha_{XY} = 0$.

Remark 10. The fundamental form α_{XY} is closed, i.e. $d\alpha_{XYZ} = 0$, moreover $\beta_{XYZ} = 0$.

The corollary below follows from Theorem 1 and Remark 10.

Corollary. *The deviations A_{XY} define an 1-dimensional central extension of $\mathfrak{w}^{\mathbb{C}}$ by $\text{Im } A / (\text{Im } A \cap \mathcal{TC}_0)$ (here $\text{Im } A \subseteq \mathcal{TC}$ is the image of the mapping $A : \Lambda^2(\mathfrak{w}^{\mathbb{C}}) \mapsto \mathcal{TC}$ and $\mathcal{TC}_0$ is the class of all operators from $\mathcal{TC}$ of zero trace) with the central charge*

$$c = -2(6h^2 - 6h + 1).$$

Otherwords, the q_R -conformal symmetries define the $\mathcal{TC}_0$ -projective representation ($\mathcal{TC}_0$ is a Lie algebra) of the Virasoro algebra $\text{vir}^{\mathbb{C}}$ with this central charge.

Theorem 2A. *For the $\mathcal{TC}$ -projective representations of the Witt algebra in the unitarizable Verma modules V_h ($h \neq \frac{1}{2} \pm \frac{\sqrt{3}}{6}$) over $\mathfrak{sl}(2, \mathbb{C})$ the kernels $\text{Ker } A$ and $\text{Ker } \alpha$ coincide, therefore, the formal series $\alpha^{XY} A_{XY}$ are defined. The series absolutely converges if $h > \frac{1}{2}$, $h \neq \frac{1}{2} + \frac{\sqrt{3}}{6}$. The unitarizable Verma modules V_h are $\mathfrak{sl}(2, \mathbb{C})$ -irreducible, therefore, the mean deviation A is proportional to the identical operator E . Precisely,*

$$\alpha = -\frac{3}{2h - 1}.$$

An estimation of the convergence and formulas for α follow from the explicit computations.

Corollary. *The $\mathcal{TC}$ -projective representations of the Witt algebra in the unitarizable Verma modules V_h ($h > \frac{1}{2}$, $h \neq \frac{1}{2} + \frac{\sqrt{3}}{6}$) over $\mathfrak{sl}(2, \mathbb{C})$ are einsteinian of the second kind with $\varrho^{(2)} = \alpha$.*

Remark 11. The central charge c and the deviation constant α obey the identity

$$\alpha^2(1 - c) = 27.$$

Theorem 2B. *For $h > \frac{1}{2}$, $h \neq \frac{1}{2} + \frac{\sqrt{3}}{6}$ the formal series for the Ricci tensor $R_{XY}^{(1)}$ converges and the $\mathcal{TC}$ -projective representations of the Witt algebra $\mathfrak{w}^{\mathbb{C}}$ in the Verma modules V_h over $\mathfrak{sl}(2, \mathbb{C})$ are einsteinian of the first kind.*

Proof. An estimation of convergence follows from the explicit formulas for q_R -conformal symmetries. If the Ricci tensor $R_{XY}^{(1)}$ exists then it is $\mathfrak{sl}(2, \mathbb{C})$ -invariant. However, any $\mathfrak{sl}(2, \mathbb{C})$ -invariant 2-form on the Witt algebra $\mathfrak{w}^{\mathbb{C}}$ is proportional to the Gelfand-Fuchs cocycle.

Remark 12A. It is not known whether the Ricci constants $\varrho^{(1)}$ and $\varrho^{(2)}$ coincide.

Remark 12B. It is not known also whether the Ricci identity [15,16]

$$R_{XY,ZV} + R_{YZ,XV} + R_{ZX,YV} = 0$$

holds for the curvature form R .

It is rather interesting to calculate some “differential geometric” characteristics of q_R -conformal symmetries such as “scalar curvature”

$$K_{ij} = \frac{R_{i,-j,j,-i}}{\alpha_{i,-i}\alpha_{j,-j}}$$

and to verify whether the other identities of Riemann geometry besides Ricci one [15,16] hold for the curvature form R .

Remark 13. The constants c and α may be written in terms of q_R instead of h as

$$c = 1 - 3q_R^{-2}, \quad \alpha = -3q_R.$$

Hence,

$$\alpha \xrightarrow{q_R \rightarrow 0} 0.$$

The central charge c is equal to 0 iff $q_R = \frac{1}{\sqrt{3}}$.

Theorem 2C. *Let put*

$$B_X = \alpha^{YZ} B_{XYZ} = \alpha^{YZ} A_{[XY],Z}$$

for $h > \frac{1}{2}$, $h \neq \frac{1}{2} + \frac{\sqrt{3}}{6}$. Then

$$B_X = \alpha T(X).$$

Proof. Note that the operators B_X form the family of tensor operators in the Verma module V_h , which transforms under $\mathfrak{sl}(2, \mathbb{C})$ just as e_k , and, hence, B_X is proportional to $T(X)$ because such family of tensor operators is unique in V_h . To calculate the multiplier one should put $X = e_0$ and to apply both B_X and $T(X)$ to the extremal vector.

Remark 14. The construction of Theorem 3C allows to define a mapping from $\text{End}(\mathfrak{w}^{\mathbb{C}})$ to $\text{End}(V_h)$, namely, the operator $Q \in \text{End}(\mathfrak{w}^{\mathbb{C}})$ is mapped to the operator

$$\hat{Q}_h = \frac{1}{\alpha} \alpha^{YZ} A_{Q(Y),Z}$$

in V_h (certainly, the convergence of series depends on Q).

It is very interesting to investigate the asymptotical properties of the mapping $Q \mapsto \hat{Q}_h$ (as a mapping of one associative algebra into another) at $h \rightarrow \frac{1}{2}$, $h \rightarrow 1$ (cf.[6]) and $h \rightarrow \infty$.

3. INFINITE DIMENSIONAL GEOMETRY OF HOMOGENEOUS KÄHLER MANIFOLDS $M = \text{Diff}_+(\mathbb{S}^1)/\mathbb{S}^1$ AND $M_1 = \text{Diff}_+(\mathbb{S}^1)/\text{PSL}(2, \mathbb{R})$

3.1. Group $\text{Diff}_+(\mathbb{S}^1)$ of diffeomorphisms of a circle $\mathbb{S}^1$ and the Virasoro-Bott group Vir. Let $\text{Diff}(\mathbb{S}^1)$ denote the group of all diffeomorphisms of the circle $\mathbb{S}^1$. The group manifold $\text{Diff}(\mathbb{S}^1)$ splits into two connected components, the subgroup $\text{Diff}_+(\mathbb{S}^1)$ and the coset $\text{Diff}_-(\mathbb{S}^1)$. The diffeomorphisms in $\text{Diff}_+(\mathbb{S}^1)$ preserve the orientation on the circle $\mathbb{S}^1$ and those in $\text{Diff}_-(\mathbb{S}^1)$ reverse it. The Lie algebra of $\text{Diff}_+(\mathbb{S}^1)$ can be identified with $\text{Vect}(\mathbb{S}^1)$.

The infinite-dimensional group Vir corresponding to the Virasoro algebra vir (more precisely, to the central extension of the Lie algebra $\text{Vect}(\mathbb{S}^1)$ defined by the Gelfand-Fuchs cocycle, whereas the Virasoro algebra vir is an extension of the real form $\mathfrak{w}$ of the Witt algebra $\mathfrak{w}^{\mathbb{C}}$) is a central extension of the group $\text{Diff}(\mathbb{S}^1)$. The corresponding 2-cocycle was calculated by R.Bott [17]:

$$c(g_1, g_2) = \int \log(g'_1 \circ g_2) \log(g'_2).$$

The group Vir is called the Virasoro-Bott group.

3.2. Flag manifold $M = \text{Diff}_+(\mathbb{S}^1)/\mathbb{S}^1$ of the Virasoro-Bott group. The flag manifold M of the Virasoro-Bott group is a homogeneous space with transformation group $\text{Diff}_+(\mathbb{S}^1)$ and isotropy group $\mathbb{S}^1$. There exist several different realizations of this manifold [18-20].

Algebraic realization. The space M can be realized as a conjugacy class in the group $\text{Diff}_+(\mathbb{S}^1)$ or in the Virasoro-Bott group Vir .

Probabilistic realization. Let P be the space of real probability measures $\mu = u(t) dt$ with smooth positive density $u(t)$ on $\mathbb{S}^1$. The group $\text{Diff}_+(\mathbb{S}^1)$ naturally acts on P by the formula

$$g : u(t) dt \mapsto u(g^{-1}(t)) dg^{-1}(t).$$

The action is transitive and the stabilizer of the point $(2\pi)^{-1} dt$ is isomorphic to $\mathbb{S}^1$, therefore, P can be identified with M .

Orbital realization. The space M can be considered as an orbit of the coadjoint representation of $\text{Diff}_+(\mathbb{S}^1)$ or Vir . Namely, the elements of the dual space vir^* of the Virasoro algebra vir are identified with the pairs $(p(t) dt^2, b)$; the coadjoint action of Vir has the form

$$K(g)(p, b) = (gp - bS(g), b),$$

where

$$S(g) = \frac{g'''}{g'} - \frac{3}{2} \left(\frac{g''}{g'} \right)^2$$

is the Schwarzian (the Schwarz derivative) and gp denotes the natural action of g on the quadratic differential p . The orbit of the point $(a \cdot dt^2, b)$ coincides with M if $a/b \neq -n^2/2$, $n = 1, 2, 3, \dots$. Therefore, a family $\omega_{a,b}$ of symplectic structures (Kirillov forms) is defined on M (cf.[21]).

Analytic realization. Let us consider the space S of univalent functions on the unit disk D_+ [22-24]. The Taylor coefficients $c_1, c_2, c_3, \dots$ in the expansion

$$f(z) = z + c_1 z^2 + c_2 z^3 + \dots + c_n z^{n+1} + \dots$$

form a coordinate system on S . The class S can be naturally identified with M via the Kirillov construction [20]. The Lie algebra $\text{Vect}(\mathbb{S}^1)$ acts on S by the formulas

$$L_v(f(z)) = if^2(z) \oint \left(\frac{wf'(w)}{f(w)} \right)^2 \frac{v(w)}{f(w) - f(z)} \frac{dw}{w}.$$

The Kirillov construction supply M by the complex structure. The symplectic structure $\omega_{a,b}$ coupled with the complex structure determines a Kähler metric $w_{a,b}$ on M . More detailed information on the infinite-dimensional geometry of the flag manifold M is contained in [2] (see also refs wherein). Note only that the curvature tensors of the Kähler connections on M were calculated in [1].

3.3. Infinite dimensional Kähler manifold $M_1 = \text{Diff}_+(\mathbb{S}^1)/\text{PSL}(2, \mathbb{R})$. The subgroup $\mathbb{S}^1$ is contained in each of the subgroups H_k , $k = 1, 2, 3, \dots$, generated by the generators ie_0 , $e_k - e_{-k}$ and $i(e_k + e_{-k})$. The subgroup H_k is isomorphic to the k -folded covering of the group $H_1 = \text{PSL}(2, \mathbb{R})$ and acts on $\mathbb{S}^1$ by the formulas

$$z \mapsto \left(\frac{az^k + b}{\bar{b}z^k + \bar{a}} \right)^{1/k} = \alpha z \left(\frac{1 + bz^{-k}}{1 + \bar{b}z^k} \right)^{1/k},$$

where $\alpha = (a/|a|)^{2/k}$ is an univalued function on H_k .

The homogeneous space $M_k = \text{Diff}_+(\mathbb{S}^1)/H_k$ is a symplectic manifold and can be identified with the orbit with $a/b = -k^2/2$ in the coadjoint representation of the Virasoro-Bott group. The manifold M_k is a quotient of M by the $\text{Diff}_+(\mathbb{S}^1)$ -invariant foliation $\mathcal{F}_k$ generated by s_k and c_k (considered as elements of a reductive basis on M [15]). Note that though M is a reductive space the manifolds M_k are not reductive. The foliations $\mathcal{F}_k$ are complex foliations, which were discovered by V.Yu.Ovsienko and O.D.Ovsienko. The foliation $\mathcal{F}_1$ is holomorphic whereas other foliations are not holomorphic. Hence, the almost complex structure on M_1 is integrable and, therefore, $M_1 = \text{Diff}_+(\mathbb{S}^1)/\text{PSL}(2, \mathbb{R})$ is a homogeneous Kähler manifold. This is not true for $k \geq 2$.

The main characteristics of the Kähler metric on M_1 were calculated in [1] (really, there were calculated their back-liftings to M). In particular, these Kähler metrics are einsteinian.

3.4. Deformed holomorphic tangent bundles on $M_1 = \text{Diff}_+(\mathbb{S}^1)/\text{PSL}(2, \mathbb{R})$, $\text{Diff}_+(\mathbb{S}^1)$ -invariant hermitean connections and their curvatures. Note that one can deform the holomorphic tangent bundle $T^{1,0}(M_1)$ on M_1 in the following way. Let us consider the principal $\text{PSL}(2, \mathbb{R})$ -bundle $\mathcal{P}$ over the homogeneous manifold M_1 and the associated holomorphic bundle $T_h(M_1)$ with the fiber isomorphic to V_h . The bundle $T_h(M_1)$ is supplied by the holomorphic action of the Witt algebra $\mathfrak{w}^\mathbb{C}$, which is integrable to the action of the group $\text{Diff}_+(\mathbb{S}^1)$. The holomorphic tangent bundle $T^{1,0}(M_1)$ is just the bundle $T_2(M_1)$. The hermitean form in V_h ($h > \frac{1}{2}$) supplies $T_h(M_1)$ by a structure of the hermitean bundle.

Now we are able to formulate the main theorem of the article.

Theorem 3A. *There exists a unique holomorphic $\mathfrak{w}^\mathbb{C}$ -invariant ($\text{Diff}_+(\mathbb{S}^1)$ -invariant) hermitean connection in $T_h(M_1)$. The Nomizu operators of the connection coincides with the generators of q_R -conformal symmetries L_k , whereas its curvature coincides with the deviations A_{XY} .*

Note that though M_1 is not reductive the Nomizu operators are correctly defined (one can use the back-lifting to the reductive space M).

Proof. One should use the definition of q_R -conformal symmetries. Indeed, L_k ($k \leq 1$) define the natural action of the corresponding generators of the Witt algebra in the fiber V_h of the bundle $T_h(M_1)$, whereas $L_{-k} = L_k^*$.

Corollary. *The deviations A_{XY} coincides with the curvature of Kähler metric on M_1 [1] at $h = 2$.*

Remark 15. The curvature form R of the $\mathcal{HS}$ -projective representation of the Witt algebra $\mathfrak{w}^\mathbb{C}$ in the Verma module V_h over the Lie algebra $\mathfrak{sl}(2, \mathbb{C})$ is just the trace of a square of the curvature of holomorphic $\mathfrak{w}^\mathbb{C}$ -invariant hermitean connection in the deformed holomorphic tangent bundle $T_h(M_1)$ over the infinite dimensional Kähler manifold $M_1 = \text{Diff}_+(\mathbb{S}^1)/\text{PSL}(2, \mathbb{R})$.

Let us reformulate Theorem 1 in the following way.

Theorem 3B. *The holomorphic $\mathfrak{w}^\mathbb{C}$ -invariant hermitean connection in the deformed holomorphic tangent bundle $T_h(M_1)$ over the infinite-dimensional Kähler manifold $M_1 = \text{Diff}_+(\mathbb{S}^1)/\text{PSL}(2, \mathbb{R})$ defines the prequantization connection [25] in*

the determinant bundle $\det T_h(M_1)$ over this manifold, which supplies $\det T_h(M_1)$ by an action of the Virasoro algebra $\text{vir}^{\mathbb{C}}$ with the central charge

$$c = -2(6h^2 - 6h + 1).$$

Remark 16. At $h = 2$ the central charge is equal to -26 as it was once calculated in [26].

CONCLUSIONS

Thus, we established a geometric interpretation of approximate ($\mathcal{HS}$ -projective or $\mathcal{TC}$ -projective) representations of the Witt algebra $\mathfrak{w}^{\mathbb{C}}$ by q_R -conformal symmetries in the Verma modules V_h over the Lie algebra $\mathfrak{sl}(2, \mathbb{C})$ and calculated some their characteristics. The generators of representations coincide with the Nomizu operators of holomorphic $\mathfrak{w}^{\mathbb{C}}$ -invariant hermitean connections in the deformed holomorphic tangent bundles $T_h(M_1)$ over the infinite-dimensional Kähler manifold $M_1 = \text{Diff}_+(\mathbb{S}^1)/\text{PSL}(2, \mathbb{R})$, whereas the deviations A_{XY} of the approximate representations coincide with the curvature operators for these connections, which supply the determinant bundle $\det T_h(M_1)$ by a structure of the prequantization bundle over M_1 . At $h = 2$ the geometric picture reduces to one considered in [1] (without any relation to approximate representations) for ordinary tangent bundles.

REFERENCES

- [1] Kirillov A.A., Juriev D.V., The Kähler geometry of the infinite dimensional homogeneous space $M = \text{Diff}_+(\mathbb{S}^1)/\text{Rot}(\mathbb{S}^1)$. *Funct. Anal. Appl.* 21(4) (1987) 284-293.
- [2] Juriev D.V., The vocabulary of geometry and harmonic analysis on the infinite-dimensional manifold $\text{Diff}_+(\mathbb{S}^1)/\mathbb{S}^1$. *Adv. Soviet Math.* 2 (1991) 233-247.
- [3] Juriev D., Quantum conformal field theory as infinite dimensional noncommutative geometry. *Russian Math. Surveys* 46(4) (1991) 135-163.
- [4] Juriev D., Infinite dimensional geometry and quantum field theory of strings. I-III. *Alg. Groups Geom.* 11(2) (1994) 145-179, *Russian J. Math. Phys.* 4(3) (1996) 287-314, *J. Geom. Phys.* 16 (1995) 275-300.
- [5] Juriev D., String field theory and quantum groups. I.: q-alg/9708009.
- [6] Juriev D., Approximate representations and Virasoro algebra: math.RT/9805001.
- [7] Zhelobenko D.P., Representations of the reductive Lie algebras [in Russian]. Moscow, Nauka, 1993.
- [8] Dixmier J., Algèbres enveloppantes. Gauthier-Villars, Paris, 1974.
- [9] Barut A., Raczka R., Theory of group representations and applications. PWN – Polish Sci. Publ. Warszawa, 1977.
- [10] Juriev D.V., Complex projective geometry and quantum projective field theory. *Theor. Math. Phys.* 101(3) (1994) 1387-1403.
- [11] Berezin F.A., Quantization in complex symmetric spaces [in Russian]. *Izvestiya AN SSSR. Ser. matem.* 39(2) (1975) 363-402.
- [12] Klimek S., Lesniewski A., Quantum Riemann surfaces. I. *Commun. Math. Phys.* 146 (1992) 103-122.
- [13] Biedenharn L., Louck J., Angular momentum in quantum mechanics. Theory and applications. *Encycl. Math. Appl.* V.8. Addison Wesley Publ. Comp. 1981.
- [14] Fuchs D.B., Cohomology of infinite-dimensional Lie algebras [in Russian]. Moscow, Nauka, 1984.
- [15] Kobayashi Sh, Nomizu K., Foundations of differential geometry. *Intersci. Publ.*, 1963/69.
- [16] Rashevskii P.K., Riemann geometry and tensor analysis [in Russian]. Moscow, 1967.
- [17] Bott R., The characteristic classes of groups of diffeomorphisms. *Enseign. Math.* 23 (1977) 209-220.

- [18] Kirillov A.A., Infinite-dimensional Lie groups, their invariants and representations. Lect. Notes Math. 970 (1982) 101-123.
- [19] Segal G., Unitary representations of some infinite-dimensional groups. Commun. Math. Phys. 80 (1981) 301-342.
- [20] Kirillov A.A., A Kähler structure on K-orbits of the group of diffeomorphisms of the circle. Funct.Anal.Appl. 21 (1987) 322-324.
- [21] Kirillov A.A., Elements of representation theory. Springer, 1976.
- [22] Goluzin G.M., Geometric theory of functions of complex variables. Amer. Math. Soc. 1968.
- [23] Duren P.L., Univalent functions. Springer, 1983.
- [24] Lehto O., Univalent functions and Teichmüller spaces. Springer, 1986.
- [25] Kostant B., Quantization and unitary representations. Lect. Notes Math. 170 (1970) 87-208.
- [26] Bowick M.J., Rajeev S.G., The holomorphic geometry of closed bosonic strings and $\text{Diff}(S^1)/S^1$. Nucl. Phys. B293 (1987) 348-384.